

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
23 January 2003 (23.01.2003)

PCT

(10) International Publication Number
WO 03/006419 A1

(51) International Patent Classification⁷: C07C 209/20,
211/63

(21) International Application Number: PCT/US02/21236

(22) International Filing Date: 2 July 2002 (02.07.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/303,971 9 July 2001 (09.07.2001) US

(71) Applicant (for all designated States except US): LONZA
INC. [US/US]; 17-17 Route 208, Fair Lawn, NJ 07410
(US).

(72) Inventor; and

(75) Inventor/Applicant (for US only): WALKER, Leigh, E.
[US/US]; 4758 Macungie Mountain Road, Macungie, PA
18062 (US).

(74) Agents: LESSLER, Jay, P. et al.; Darby & Darby P.C.,
805 Third Avenue, New York, NY 10022-7513 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, EC, EE, ES, FI, GB, GD, GE, HR, HU,
ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS,
LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO,
NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ,
TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK,
TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

WO 03/006419 A1

(54) Title: IN SITU PROCESS FOR PREPARING QUATERNARY AMMONIUM BICARBONATES AND QUATERNARY AM-
MONIUM CARBONATES

(57) Abstract: Applicant has discovered an *in situ* method of preparing quaternary ammonium methocarbonate salts and quaternary ammonium alkylcarbonate salts in high yield from tertiary amines, methanol, and at least one of a cyclic carbonate, an aliphatic polyester (such as a polycarbonate), or an ester (such as a carbonate ester), and their subsequent conversion to quaternary ammonium bicarbonates, quaternary ammonium carbonates or both in a one-pot reaction. According to one embodiment of the invention, the method includes reacting an amine and methanol with at least one of a cyclic carbonate and an aliphatic polyester to yield a quaternary ammonium methocarbonate. This method does not produce or require the handling of corrosive quaternary ammonium hydroxides. Another embodiment is a method of preparing quaternary ammonium alkylcarbonate salts by reacting tertiary amines, methanol, and an ester. The quaternary ammonium methocarbonate or alkylcarbonate can be converted to the corresponding bicarbonate, carbonate, or mixture thereof by methods known in the art.

**IN SITU PROCESS FOR PREPARING QUATERNARY AMMONIUM
BICARBONATES AND QUATERNARY AMMONIUM CARBONATES**

This application claims the benefit of U.S. Provisional Application No. 60/303,971, filed July 9, 2001, which is hereby incorporated by reference.

5 **Field of the Invention**

This invention relates to an *in situ* method of preparing quaternary ammonium alkylcarbonates (such as quaternary ammonium methocarbonates), quaternary ammonium bicarbonates, and quaternary ammonium carbonates from corresponding tertiary amines.

10 **Background of the Invention**

Quaternary ammonium compounds, such as didecyldimethyl ammonium carbonate and didecyldimethyl ammonium chloride, are known to have antimicrobial activity. See, for example, U.S. Patent Nos. 5,523,487, 5,833,741, and 6,080,789. Quaternary ammonium compounds have been found to be particularly useful as wood
15 preservatives. However, quaternary ammonium chlorides have been found to leach rapidly in soil (Nicholas *et al.*, *Forest Prod. J.*, 41:41 (1991)). Consequently, a metal coupler, such as a copper salt, is frequently added to the quaternary ammonium chlorides to prevent leaching.

Quaternary ammonium carbonates, on the other hand, have better leaching
20 resistance and do not require the use of a metal coupler. As a result, there is an increasing demand in the preservative market for quaternary ammonium carbonates.

U.S. Patent No. 5,438,034 discloses a process for preparing quaternary ammonium carbonates. The process includes reacting a quaternary ammonium chloride with a metal hydroxide to form a quaternary ammonium hydroxide and then reacting the quaternary ammonium hydroxide with carbon dioxide to yield the quaternary ammonium carbonate. The quaternary ammonium hydroxide, however, is very corrosive. In addition, metal chloride produced as a byproduct in the first reaction must be filtered out of the reaction product, a step which increases the cost and decreases the efficiency of the process. Thus, an alternative method to produce quaternary ammonium carbonates is desirable.

Wernitz, U.S. Patent No. 2,635,100, discloses a process for preparing quaternary ammonium carbonates by reacting a trialkylamine with a dialkyl carbonate, such as dimethyl carbonate and ethylene carbonate, preferably in the presence of an alcohol. Wernitz reported that the reaction of tertiary amines and dimethyl carbonate yielded quaternary ammonium methocarbonates. Wernitz also reported that when ethylene carbonate was reacted with triethylamine and methanol, the cyclic carbonate of triethyl-2-hydroxyethylammonium hydroxide was formed. The solvent, unreacted amine, and cyclic ester were removed by distillation. Many dialkyl carbonates of carbonic acid, such as dimethyl carbonate, are expensive and, therefore, significantly increase the cost of preparing quaternary ammonium carbonates by this process.

Dimethyl carbonate is commercially available, and methods of its synthesis are well known in the art. Typically cyclic carbonates, *e.g.*, ethylene and propylene carbonate, are converted to dimethyl carbonate and a glycol in the presence of methanol or other alcohol and catalyst. Romano *et al.*, U.S. Patent No. 4,062,884, disclose a process for preparing dialkylcarbonates by reacting an alcohol with a cyclic carbonate in the presence of an organic base, such as a tertiary aliphatic amine. Romano *et al.* describe the reaction of a methanol/ethylene carbonate/triethylamine mixture. Continuous distillation of the methanol-dimethylcarbonate azeotrope over 3 hours resulted in almost complete conversion of ethylene carbonate to ethylene glycol and dimethyl carbonate. Romano *et al.* further teach that the organic base which catalyzes the reaction can be totally recovered from the reaction vessel by simple distillation.

There is a continuing need for cheaper and more efficient methods for preparing quaternary ammonium carbonates. A one step, *in situ* method of preparing quaternary ammonium carbonates would advantageously meet these needs.

Summary of the Invention

5 Applicant has discovered an *in situ* method of preparing quaternary ammonium methocarbonate salts and quaternary ammonium alkylcarbonate salts in high yield from tertiary amines, methanol, and at least one of a cyclic carbonate, an aliphatic polyester, and an ester, and their subsequent conversion to quaternary ammonium bicarbonates, quaternary ammonium carbonates or both in a one-pot reaction. According to one embodiment of the invention, the method includes reacting an amine and methanol with at least one of a cyclic carbonate, an aliphatic polyester (such as a polycarbonate), or an ester (such as a carbonate ester) to yield a quaternary ammonium methocarbonate. This method does not produce or require the handling of corrosive quaternary ammonium hydroxides. Furthermore, this method advantageously produces glycols as byproducts.

10 Glycols are frequently added to solutions containing quaternary ammonium carbonates and quaternary ammonium bicarbonates to raise their flashpoint and as an anti-freeze.

 Another embodiment is a method of preparing quaternary ammonium alkylcarbonate salts by reacting tertiary amines, methanol, and an ester.

 The present invention also provides a method of preparing a quaternary ammonium bicarbonate, quaternary ammonium carbonate, or mixture thereof by (a) preparing a quaternary ammonium methocarbonate or a quaternary ammonium alkylcarbonate by one of the aforementioned methods, and (b) converting the quaternary ammonium methocarbonate or quaternary ammonium alkylcarbonate to the corresponding quaternary ammonium bicarbonate, quaternary ammonium carbonate, or mixture thereof.

20

Detailed Description of the Invention


25 Applicant has discovered an *in situ* method of preparing quaternary ammonium methocarbonate and alkylcarbonate salts in high yield from tertiary amines, methanol, and at least one of a cyclic carbonate, an aliphatic polyester (such as a polycarbonate), and/or an ester (such as a carbonate ester), and their subsequent

conversion to quaternary ammonium bicarbonates, quaternary ammonium carbonates, or mixtures thereof in a one-pot reaction.

The term "alkyl" as used herein includes straight and branched alkyl substituents. An "alkyl" group is a saturated hydrocarbon.

5 Preparation of the Quaternary Ammonium Methocarbonate

The method of the present invention can prepare quaternary ammonium methocarbonates having the formula



where R^1 , R^2 , and R^3 are independently C_1 - C_{30} alkyl. Preferably, R^1 , R^2 , and R^3 are
 10 independently C_1 - C_{20} alkyl and more preferably are independently C_1 - C_{16} alkyl. Most preferably, R^1 is methyl.

Also, at least one of R^1 , R^2 , and R^3 is preferably a C_8 - C_{30} alkyl or a C_8 - C_{20} alkyl. According to one embodiment, R^1 and R^2 are independently C_1 - C_{20} alkyl and more preferably are independently C_1 - C_{16} alkyl and R^3 is C_8 - C_{20} alkyl and more preferably C_8 -
 15 C_{16} alkyl.

According to a preferred embodiment, R^2 is C_1 - C_{20} alkyl. According to another embodiment, R^2 is methyl. According to yet another embodiment, R^2 is a C_8 - C_{12} alkyl and more preferably a C_{10} alkyl.

According to a preferred embodiment, R^3 is a C_8 - C_{12} alkyl and more
 20 preferably a C_{10} alkyl.


According to another preferred embodiment, R^1 is methyl and R^2 and R^3 are independently C_8 - C_{20} alkyl. More preferably, R^2 and R^3 are independently C_8 - C_{12} alkyl. According to a more preferred embodiment, R^2 and R^3 are C_{10} alkyl.

According to yet another preferred embodiment, R^1 and R^2 are methyl and R^3 is a C_8 - C_{20} alkyl. According to one embodiment, R^3 is a C_{10} - C_{18} alkyl and more preferably is a C_{12} or C_{18} alkyl. According to another embodiment, R^3 is C_8 - C_{12} alkyl and more preferably is a C_{10} alkyl.

5 Representative quaternary ammonium methocarbonates include, but are not limited to, didecyltrimethyl ammonium methocarbonate, dodecyltrimethyl ammonium methocarbonate, dioctyltrimethyl ammonium methocarbonate, octadecyltrimethyl ammonium methocarbonate, dioctadecyltrimethyl ammonium methocarbonate, trioctylmethyl ammonium methocarbonate, and any combination of any of the foregoing.


10 The quaternary ammonium methocarbonate is prepared by reacting an amine and methanol with at least one of a cyclic carbonate, an aliphatic polyester (such as a polycarbonate), or an ester (such as a carbonate ester). Suitable amines include, but are not limited to, those having the formula $NR^1R^2R^3$, wherein R^1 , R^2 , and R^3 are defined as above. Preferred amines include, but are not limited to, didecylmethylamine, 15 dodecylmethylamine, dioctylmethylamine, octadecylmethylamine, dioctadecylmethylamine, trioctylamine, and any combination of any of the foregoing.

Suitable cyclic carbonates include, but are not limited to, those having the formula



20 where R^4 is hydrogen or C_1 - C_4 alkyl and n is an integer from 1 to 10. Preferably, R^4 is hydrogen or methyl. Preferred cyclic carbonates include, but are not limited to, ethylene carbonate, propylene carbonate, and mixtures thereof.

Suitable aliphatic polyester include, but are not limited to, those having the formula




where R^5 , R^6 , R^7 , and R^8 are independently hydrogen or C_1 - C_{10} alkyl and m is an integer from 1 to 1200. According to one embodiment, R^5 , R^6 , R^7 , and R^8 are independently hydrogen or C_1 - C_4 alkyl. Preferably, R^5 , R^6 , R^7 , and R^8 are independently hydrogen or methyl. According to a preferred embodiment, R^5 is methyl and R^6 , R^7 , and R^8 are hydrogen. Preferably, m ranges from 1 to 100.

Generally, the reaction is performed with a molar excess of methanol and cyclic carbonate, aliphatic polyester, or mixture thereof with respect to amine, *i.e.*, the molar ratios of methanol and the cyclic carbonate, aliphatic polyester, or mixture thereof to amine are greater than 1. The molar ratio of amine to cyclic carbonate, aliphatic polyester, or mixture thereof preferably ranges from about 1:1 to about 1:10 and more preferably ranges from about 1:1.1 to about 1:1.3. The molar ratio of amine to methanol broadly ranges from about 1:2 to about 1:20 and preferably ranges from about 1:3 to about 1:10.

The reaction is typically performed at from about 120 to about 160° C, preferably from about 120 to about 150° C, and more preferably from about 120 to about 140° C. The reaction may be performed at a pressure ranging from about 60 to about 200 psi. Preferably, the reaction is performed at a pressure ranging from about 120 to about 150 psi.


Generally, the reaction is performed for from about 3 to about 40 hours and preferably from about 5 to 30 hours.

The reaction step yielding the quaternary ammonium methocarbonate also produces glycols, such as ethylene glycol and propylene glycol, as byproducts. For example, when the cyclic carbonate is propylene carbonate, the reaction is as follows:



where R^1 , R^2 , and R^3 are defined as above.

Without being bound by any theory, the inventor believes that the propylene carbonate first reacts with methanol to form dimethyl carbonate and propylene glycol as shown below.



The amine $\text{NR}^1\text{R}^2\text{R}^3$ catalyzes this first reaction. The inventor further hypothesizes that the dimethyl carbonate formed reacts with the amine to yield the quaternary ammonium methocarbonate. Therefore, the reaction yields quaternary ammonium methocarbonate and propylene glycol.

Dimethyl carbonate may also be added to the reaction mixture to improve the kinetics of the reaction, *i.e.*, to accelerate the reaction. The dimethyl carbonate also functions to balance the ratio of quaternary ammonium bicarbonate and carbonate to glycol in the product. The molar ratio of amine to dimethyl carbonate (before the reaction) preferably ranges from about 2:1 to about 1:3 and more preferably ranges from about 1.25:1 to about 1:1.25.

The inventor has found that the molar ratio of total carbonate sources (*i.e.* total cyclic carbonate and aliphatic polyester) to amine in the reaction is preferably in the range of from about 1:1 to 5:1, more preferably 1.25:1 to 2.5:1, and most preferably 1:1.5 to 1:2.


Thus, the present reaction in a preferred embodiment uses a molar excess of methanol and dimethyl carbonate relative to amine to substantially completely convert the amine to quaternary ammonium methocarbonate. The cyclic carbonate is preferably substantially completely converted to glycol and dimethyl carbonate. The glycol can remain in the product.

The molar ratio of cyclic carbonate to amine can be varied to obtain the desired weight ratio of glycol to quaternary ammonium methocarbonate in the reaction product. While methanol in the final product can be easily distilled off, removal of glycols is much more costly. According to one embodiment, the molar ratio of cyclic carbonate to amine is slightly greater than 1, *e.g.*, from about 1:1.1 to about 1:1.5 or from about 1:1.1 to about 1:1.3. By keeping the molar ratio near 1, the weight ratio of propylene glycol to quaternary ammonium methocarbonate, such as didecyldimethyl ammonium methocarbonate, in the final product generally ranges from about 1:3 to about 1:7 and is preferably about 1:5.

After the formation of the quaternary ammonium methocarbonate, excess methanol and dimethyl carbonate may be removed and recovered by simple distillation. The quaternary ammonium methocarbonate may be isolated and purified by methods known in the art.

Preparation of the Quaternary Ammonium Alkylcarbonate

According to another embodiment of the present invention, quaternary ammonium alkylcarbonates having the formula




5 where R^1 , R^2 , and R^3 are defined as above and R^9 is a C_1 - C_{10} alkyl. According to one preferred embodiment, R^9 is a C_1 - C_4 alkyl. More preferably, R^9 is methyl, ethyl, or propyl.

Representative quaternary ammonium alkylcarbonates include, but are not limited to, didecyldimethyl ammonium ethylcarbonate, dodecyltrimethyl ammonium ethylcarbonate, dioctyldimethyl ammonium ethylcarbonate, octadecyltrimethyl ammonium ethylcarbonate, dioctadecyldimethyl ammonium ethylcarbonate, trioctylmethyl ammonium ethylcarbonate, and any combination of any of the foregoing.

The quaternary ammonium alkylcarbonate is prepared by reacting an amine and methanol with an ester. Suitable amines include, but are not limited to, those described above.

15 Suitable esters include, but are not limited to, those having the formula



wherein R^9 is defined as above and R^{10} is a C_1 - C_{10} alkyl. According to one embodiment, R^{10} is a C_1 - C_4 alkyl, such as methyl, ethyl, or propyl. A preferred ester is diethyl carbonate (*i.e.* where R^9 and R^{10} are ethyl)


20 Generally, the reaction is performed with a molar excess of methanol and ester with respect to amine, *i.e.*, the molar ratios of methanol and ester to amine are greater than 1. The molar ratio of amine to ester preferably ranges from about 1:1 to about 1:10 and more preferably ranges from about 1:1.1 to about 1:1.3. The molar ratio of amine to

methanol broadly ranges from about 1:2 to about 1:20 and preferably ranges from about 1:3 to about 1:10.

The reaction conditions are generally the same as those described for the preparation of the methocarbonate.

5

The reaction step yielding the quaternary ammonium alkylcarbonate also produces alkanols having the formula R^9OH , $R^{10}OH$, or mixtures thereof. For example, when the ester is cyclic carbonate is propylene carbonate, the reaction is as follows:



where R^1 , R^2 , R^3 , R^9 , and R^{10} are defined as above.

10

Without being bound by any theory, the inventor believes that the ester first reacts with methanol to form a methyl ester of the formula $CH_3OC(O)OR^9$, $CH_3OC(O)OR^{10}$, or a mixture thereof and R^9OH , $R^{10}OH$, or a mixture thereof. The amine $NR^1R^2R^3$ catalyzes this first reaction. The inventor further hypothesizes that the methyl

ester formed reacts with the amine to yield the quaternary ammonium alkylcarbonate. Therefore, the reaction yields quaternary ammonium alkylcarbonate and alkanol (R^9OH , $R^{10}OH$, or a mixture thereof). The alkylcarbonate anion of the quaternary ammonium alkyl carbonate can be $[OC(O)OR^9]$, $[OC(O)OR^{10}]$, or a mixture thereof.

5 Alkyl methyl carbonate of the formula of the formula $R^9OC(O)OCH_3$ or $R^{10}OC(O)OCH_3$ may also be added to the reaction mixture to improve the kinetics of the reaction, *i.e.*, to accelerate the reaction. The molar ratio of amine to alkyl methyl carbonate (before the reaction) preferably ranges from about 2:1 to about 1:3 and more preferably ranges from about 1.25:1 to about 1:1.25.

10 The inventor has found that the molar ratio of total carbonate sources (*i.e.* total ester) to amine in the reaction is preferably in the range of from about 1:1 to 5:1, more preferably 1.25:1 to 2.5:1, and most preferably 1:1.5 to 1:2.

Thus, the present reaction in a preferred embodiment uses a molar excess of methanol and alkyl methyl carbonate relative to amine to substantially completely convert the amine to quaternary ammonium alkylcarbonate. The ester is substantially completely converted to alkanol and alkyl methyl carbonate. The alkanol remains in the product.

15 After the formation of the quaternary ammonium alkylcarbonate, excess methanol and alkyl methyl carbonate may be removed and recovered by simple distillation. The quaternary ammonium alkylcarbonate may be isolated and purified by methods known in the art.

Conversion of the Quaternary Ammonium Methocarbonate or
Quaternary Ammonium Alkylcarbonate to
the Corresponding Quaternary Ammonium Bicarbonate

25 The quaternary ammonium methocarbonate or alkylcarbonate produced by the method of the present invention can be converted to a corresponding bicarbonate, carbonate, or mixture thereof by methods known in the art, such as hydrolysis and other exchange reactions (e.g., de-hydrolysis). For example, the methocarbonate or alkylcarbonate may be stirred with water at ambient conditions to effect the reaction (hydrolysis) to the corresponding bicarbonate and an alkanol (methanol in the case of the methocarbonate and R^9OH , $R^{10}OH$, or a mixture thereof in the case of the alkylcarbonate).


Water may then be added to distill out the water and any residual methanol or methanol or other alkanol formed when the methocarbonate or alkylcarbonate hydrolyzes to the bicarbonate. The distillation may be done at atmospheric or reduced pressures by methods standard in the art.

5 The bicarbonate can be converted to the carbonate by any method in the art. For instance, the bicarbonate can be heated (e.g. in water) to yield the corresponding carbonate, carbon dioxide, and water.


 If methanol is distilled out of the bicarbonate solution, the heat may cause some or all of the bicarbonate to be converted to the corresponding carbonate.

10 The methanol and alkanol in the product can be recovered by any method known in the art, such as distillation as discussed above, during or after the reaction. The quaternary ammonium bicarbonate may be isolated and purified by methods known in the art.

 The method of the present invention can produce quaternary ammonium
15 bicarbonates having the formula



and quaternary ammonium carbonates having the formula



20 wherein R^1 , R^2 , and R^3 are defined as above.

 According to a preferred embodiment, the conversion results in the formation of a mixture containing from about 70 to about 90% by weight of quaternary ammonium bicarbonate and from about 10 to about 30% by weight of quaternary

ammonium carbonate, based upon 100% total weight of quaternary ammonium bicarbonate and quaternary ammonium carbonate.

Mixing, adding, and reacting steps in the present invention can be accomplished by conventional means known to those of ordinary skill in the art. The order of addition of reactants or solvent in any individual step does not affect the process. Reactants and/or solvent can be added sequentially or simultaneously in any suitable reaction vessel. Importantly, the method of the present invention is suitable for commercial scale production techniques and equipment, yet convenient for small scale work.

The following examples illustrate the invention without limitation. All parts and percentages are given by weight unless otherwise indicated.

A. Measurement of the Conversion of an Amine to the Corresponding Quaternary Ammonium Methocarbonate

The amount of amine converted to the corresponding quaternary ammonium methocarbonate was determined as follows. Gas chromatography was performed on a given sample after formation of the quaternary ammonium methocarbonate. Gas chromatography measurements of the solution were performed with a Hewlett Packard Model 5890 Series II chromatograph equipped with a Hewlett Packard 7673 GC/SFC auto-injector which injects 1 μ l of solution into the injector at 300° C. The column temperature starts at 100° C with a 5 minute hold followed by ramping at 8° C per minute to 300° C, followed by another 5 minute hold.

In the gas chromatography injector, the quaternary ammonium methocarbonate decomposes. For example, didecyldimethyl ammonium methocarbonate decomposes to produce predominantly decene and decyldimethylamine and small amounts of didecylmethylamine. In a pure solution of didecyldimethyl ammonium methocarbonate, the didecylmethylamine peak contributes 13% of the areas of the total chromatogram. The percentage of didecylmethylamine and propylene carbonate converted to didecyldimethyl ammonium methocarbonate was calculated by following the decrease in area for the didecylmethylamine peak from 100% initially to 13% representing 100% conversion. The amount of other amines converted can be determined by similar methods.

A two phase titration with a standard sodium lauryl sulfate titrant and bromophenol blue indicator was performed to confirm the formation of quaternary ammonium compounds.

B. Preparation of Quaternary Ammonium Methocarbonate

5 The following examples were performed in a 1 liter stainless steel Parr Model 4520 reactor equipped with a magnetic stirrer and a turbine type impeller. The temperature of the reactor was controlled to within $\pm 1^\circ \text{C}$ by a Parr Model 4843 controller with an electrical external heater and internal cooling by a water filled coil. An Ashcroft pressure gauge capable of measuring pressures ranging from 0 to 200 psi was attached to
10 the reactor. A small bleed valve was kept partially open until the contents of the reactor reached 60°C to facilitate removal of residual air in the reactor.

Example 1

The reactor was charged with 295 g (0.95 moles) didecylmethylamine (92% (w/w) pure and 8% (w/w) tridecylamine), 153 g (1.5 moles) propylene carbonate, and 177
15 g (5.7 moles) of methanol. The mixture was heated to 140°C with stirring for 28 hours. Didecyldimethyl ammonium methocarbonate was produced.

Initially, the solution consisted of two phases. The top layer was essentially pure didecylmethyl amine. After 15 hours, the solution was a single phase. When a sample of the solution after 15 hours of heating was allowed to cool to room temperature,
20 it became hazy and separated into two phases.

According to gas chromatography, about 90 to 95% of the didecylmethylamine was converted to didecyldimethyl ammonium methocarbonate after 15 hours of heating and about 99% of the didecylmethylamine was converted to didecyldimethyl ammonium methocarbonate after 28 hours of heating.

25 Example 2

The reactor was charged with didecylmethylamine, propylene carbonate, and methanol at a molar ratio of 1:1.2:6.5. The mixture was heated to 130°C for 16 hours. Didecyldimethyl ammonium methocarbonate was produced. Initially, the solution

consisted of two phases. After heating, the solution remained in two phases. The top layer was high in tridecylamine.

According to gas chromatography, about 75-80% of the didecylmethylamine was converted to didecylmethyl ammonium methocarbonate after the 16 hours of heating.

5

Example 3

A reactor was charged with didecylmethylamine, propylene carbonate, and methanol at a molar ratio of 1:1.6:10. The mixture was heated to 130° C for 20 hours. Didecylmethyl ammonium methocarbonate was produced. Initially, the solution consisted of two phases. After 16 hours of heating, the solution consisted of one phase.

10

According to gas chromatography, about 85-90% of the didecylmethylamine was converted to didecylmethyl ammonium methocarbonate after the 20 hours of heating.

Example 4

The reactor was charged with didecylmethylamine (100% pure), propylene carbonate, and methanol at a molar ratio of 1:2:10. The mixture was heated to 140° C for 23 hours. Didecylmethyl ammonium methocarbonate was produced. Initially, the solution consisted of two phases. After 6 hours of heating, the solution consisted of one phase.

15

According to gas chromatography, about 97-100% of the didecylmethylamine was converted to didecylmethyl ammonium methocarbonate after the 23 hours of heating.

20

Example 5

The reactor was charged with 233 g (0.75 moles) didecylmethylamine (96% (w/w) pure and 4% (w/w) tridecylamine), 76.5 g (0.75 moles) propylene carbonate, 67.5 g (0.75 moles) dimethyl carbonate, and 233 g (7.5 moles) methanol. The mixture was heated to 140° C for 8.5 hours. Didecyl-dimethyl ammonium methocarbonate was

25

produced. Initially, the solution consisted of two phases. After 2.5 hours, the solution consisted of one phase.

According to gas chromatography, about 75, 95, and 99% of the didecylmethylamine was converted to didecyldimethyl ammonium methocarbonate after 2.5, 5.5, and 8.5 hours of heating.

Example 6

The reactor was charged with didecylmethylamine, propylene carbonate, methanol, and dimethyl carbonate at a molar ratio of 1:1:4:0.4. The mixture was heated to 150° C for 20 hours. Didecyldimethyl ammonium methocarbonate was produced. Initially, the solution consisted of two phases. After the 20 hours of heating, the solution consisted of two phases.

According to gas chromatography, about 75% of the didecylmethylamine was converted to didecyldimethyl ammonium methocarbonate after the 20 hours of heating.

Example 7

The reactor was charged with didecylmethylamine, propylene carbonate, methanol, and dimethyl carbonate at a molar ratio of 1:1:10:1. The mixture was heated to 130° C for 21 hours. Didecyldimethyl ammonium methocarbonate was produced. Initially, the solution consisted of two phases. After the 21 hours of heating, the solution consisted of one phase.

According to gas chromatography, about 97% of the didecylmethylamine was converted to didecyldimethyl ammonium methocarbonate after the 21 hours of heating.

Example 8

The reactor was charged with didecylmethylamine, propylene carbonate, methanol, and dimethyl carbonate at a molar ratio of 1:1:7:0.7. The mixture was heated to 140° C for 15 hours. Didecyldimethyl ammonium methocarbonate was produced.

Initially, the solution consisted of two phases. After 6 hours of heating, the solution consisted of one phase.

According to gas chromatography, about 98% of the didecylmethylamine was converted to didecyldimethyl ammonium methocarbonate after the 15 hours of heating.

Example 9

The reactor was charged with didecylmethylamine, propylene carbonate, methanol, and dimethyl carbonate at a molar ratio of 1:1:4:1. The mixture was heated to 150° C for 17 hours. Didecyldimethyl ammonium methocarbonate was produced.

Initially, the solution consisted of two phases. After the 17 hours of heating, the solution consisted of one phase.

According to gas chromatography, about 85% of the didecylmethylamine was converted to didecyldimethyl ammonium methocarbonate after 17 hours of heating.

Example 10

The reactor was charged with didecylmethylamine, propylene carbonate, methanol, and dimethyl carbonate at a molar ratio of 1:1:7:0.7. The mixture was heated to 150° C for 16 hours. Didecyldimethyl ammonium methocarbonate was produced.

Initially, the solution consisted of two phases. After 3 hours of heating, the solution consisted of one phase.

According to gas chromatography, about 98% of the didecylmethylamine was converted to didecyldimethyl ammonium methocarbonate after the 16 hours of heating.

Example 11

The reactor was charged with didecylmethylamine, propylene carbonate, methanol, and dimethyl carbonate at a molar ratio of 1:1:10:1. The mixture was heated to 120° C for 28 hours. Didecyldimethyl ammonium methocarbonate was produced.

Initially, the solution consisted of two phases. After 10 hours of heating, the solution consisted of one phase.

According to gas chromatography, about 98-100% of the didecylmethylamine was converted to didecyldimethyl ammonium methocarbonate after the 28 hours of heating.

Example 12

5 The reactor was charged with didecylmethylamine, propylene carbonate, methanol, and dimethyl carbonate at a molar ratio of 1:1:10:1. The mixture was heated to 140° C for 15 hours. Didecyldimethyl ammonium methocarbonate was produced.

According to gas chromatography, 95-98% of the didecylmethylamine was converted to didecyldimethyl ammonium methocarbonate after the 15 hours of heating.

10

Example 13

The reactor was charged with dodecyldimethylamine, propylene carbonate, and methanol at a molar ratio of 1:2:10. The mixture was heated to 140° C for 6 hours. Dodecyltrimethyl ammonium methocarbonate was produced.

15 According to gas chromatography, 96% and 100% of the dodecyldimethylamine, respectively, was converted to dodecyltrimethyl ammonium methocarbonate after 3 and 6 hours of heating.

Example 14

20 The reactor was charged with dodecyldimethylamine, propylene carbonate, dimethyl carbonate, and methanol at a molar ratio of 1:1:1:10. The mixture was heated to 140° C for 3 hours. Dodecyltrimethyl ammonium methocarbonate was produced.

According to gas chromatography, 100% of the dodecyldimethylamine was converted to dodecyltrimethyl ammonium methocarbonate after the 3 hours of heating.

Example 15

25 The reactor was charged with dioctylmethylamine, propylene carbonate, and methanol at a molar ratio of 1:2:10. The mixture was heated to 140° C for 14 hours. Dioctyldimethyl ammonium methocarbonate was produced. Initially, the solution consisted of two phases. After 5.5 hours of heating, the solution consisted of one phase.

According to gas chromatography, 85% and 98% of the dioctylmethylamine, respectively, was converted to dioctyldimethyl ammonium methocarbonate after 5.5 and 14 hours of heating.

Example 16

5 The reactor was charged with octadecyldimethylamine, propylene carbonate, and methanol at a molar ratio of 1:2:10. The mixture was heated to 140° C for 5 hours. Octadecyltrimethyl ammonium methocarbonate was produced.

According to gas chromatography, 97% of the octadecyldimethylamine was converted to octadecyltrimethyl ammonium methocarbonate after the 5 hours of heating.

10

Example 17

The reactor was charged with dioctadecylmethylamine, propylene carbonate, and methanol at a molar ratio of 1:2:10. The mixture was heated to 140° C for 26 hours. Dioctadecyldimethyl ammonium methocarbonate was produced.

15 According to gas chromatography, 80% of the dioctadecylmethylamine was converted to dioctadecyldimethyl ammonium methocarbonate.

Example 18

20 The reactor was charged with trioctylamine, propylene carbonate, and methanol at a molar ratio of 1:2:10. The mixture was heated to 140° C for 38 hours. Trioctylmethyl ammonium methocarbonate was produced. Initially, the solution consisted of two phases. After 38 hours of heating, the solution still consisted of two phase.

According to gas chromatography, 85% of the trioctylamine was converted to trioctylmethyl ammonium methocarbonate after the 38 hours of heating.

Example 19

25 The reactor was charged with didecylmethylamine, ethylene carbonate, and methanol at a molar ratio of 1:2:10. The mixture was heated to 140° C for 26 hours. Didecylmethyl ammonium methocarbonate was produced. Initially, the solution

consisted of two phases. After 6 hours of heating, the solution consisted of one phase. Furthermore, after 26 hours the solution was dark.

According to gas chromatography, 85% of the didecylmethylamine was converted to didecyldimethyl ammonium methocarbonate after the 26 hours of heating.

5 C. Preparation of Quaternary Ammonium Bicarbonate

Example 20

The mixture from Example 1 was cooled and transferred to a round bottom flask equipped with a Vigreux column having a distillation head condenser receiving flask cooled in a dry ice acetone bath and a connection to a vacuum system. The flask was
10 placed in an oil bath and heated slowly as the vacuum was dropped to 0.2 atm. Distillation continued over a 3 hour period as the bath temperature was slowly raised to 75°C. A total of 165 grams of liquid was collected. Using a density curve based on known mixtures of dimethyl carbonate (density 1.07g/cc) and methanol (density 0.79g/cc), it was estimated that the mixture from Example 1 contained 20% by weight of dimethyl carbonate. 400
15 grams of water were added to the flask and the flask was placed in the oil bath. The vacuum was slowly applied to distill out water (about 250 grams total) at a temperature of 55-75°C over a period of about 5 hours until a sample of the distillate had a density of greater than 0.98g/cc. The product contained about 600 grams of combined quaternary ammonium bicarbonate and carbonate (60%), propylene glycol and water and less than 1%
20 methanol. An additional 110 grams of water was added to dilute the mixture to yield a product containing 50% by weight of combined quaternary ammonium bicarbonate and carbonate, 16% by weight of propylene glycol, and 34% by weight of water.

Example 21

The methocarbonate in the mixture obtained in Example 7 was converted to
25 the corresponding bicarbonate and carbonate by the procedure described in Example 20. The resulting mixture contained 50% by weight of combined quaternary ammonium bicarbonate and carbonate, 10% by weight of propylene glycol, and 40% by weight of water.

Example 22

A 50 gallon reactor, equipped with a stripping condenser and a 30 gallon receiver, was charged with 116.4 lbs (0.374 lb moles) of didecylmethyl amine, 38.2 lbs (0.374 lb moles) of propylene carbonate, 33.7 lbs (0.374 lb moles) of dimethyl carbonate, and 119.7 g (3.740 lb moles) of methanol. The reactor was heated to and maintained at 140° C for 6-9 hours. The reactor was maintained at 150 psig. After 6 hours at 140° C, the reactor was sampled to determine the percentage of quaternary ammonium compound and free amine in the reactor. The reaction mixture was periodically sampled until the ratio of quaternary ammonium compound to unreacted amine was greater than 97.5:2.5. The quaternary ammonium compound was identified by NMR as didecyldimethylammonium methocarbonate.

Excess methanol and dimethyl carbonate was removed by atmospheric stripping at 100° C until no more distillate was collected. The reactor was cooled to 60° C. The content of the stripping condenser and receiver was drained into a 55 gallon drum. Full glycol cooling was applied to the stripping condenser and receiver and full vacuum (~4 mm Hg) was applied to the reactor system to further strip off any remaining dimethylcarbonate and methanol. When no more distillate was collected, the reactor was vented with nitrogen to atmospheric pressure. The content of the receiver was drained into the 55 gallon drum. A total of 129 lbs of dimethylcarbonate and methanol was collected in the drum.

To the content of the reactor at 60° C was added 109.6 lbs of water. The reactor was heated and maintained at 80° C for 2-3 hours. The didecyldimethylammonium methocarbonate was hydrolyzed to didecyldimethylammonium bicarbonate/carbonate. Additional water was added so that the final concentration of the quaternary ammonium compound was about 50-52%, the propylene glycol was about 9-11%, and methanol was less than 3.5%.


* * *

All patents, applications, articles, publications, and test methods mentioned above are hereby incorporated by reference.

Many variations of the present invention will suggest themselves to those skilled in the art in light of the above detailed description. Such obvious variations are within the full intended scope of the appended claims.


IN THE CLAIMS:

1. A method of preparing a quaternary ammonium methocarbonate having the formula




wherein R^1 and R^2 are independently C_1 - C_{30} alkyl and R^3 is a C_8 - C_{30} alkyl, the method comprising reacting

- (a) an amine having the formula $NR^1R^2R^3$;
 (b) (i) a cyclic carbonate having the formula



wherein R^4 is hydrogen or C_1 - C_4 alkyl and n is an integer from 1 to 10,

- (ii) an aliphatic polyester having the formula



1 wherein R⁵, R⁶, R⁷, and R⁸ are independently hydrogen or C₁-C₁₀ alkyl and m is an integer
2 from 1 to 1200, or

3 (iii) a mixture thereof; and

4 (c) methanol

5 to form the methocarbonate.

1 2. The method of claim 1, wherein R¹ and R² are independently C₁-C₂₀ alkyl
2 and R³ is a C₈-C₂₀ alkyl.

1 3. The method of claim 1, wherein R¹ and R² are independently C₁-C₁₆ alkyl
2 and R³ is a C₈-C₁₆ alkyl.

1 4. The method of claim 3, wherein R¹ is methyl.

1 5. The method of claim 1, wherein R² is C₁-C₂₀ alkyl.

1 6. The method of claim 5, wherein R² is methyl.

1 7. The method of claim 5, wherein R² is C₈-C₁₂ alkyl.

1 8. The method of claim 7, wherein R² is C₁₀ alkyl.

1 9. The method of claim 1, wherein R³ is C₈-C₂₀ alkyl.

1 10. The method of claim 9, wherein R³ is C₈-C₁₂ alkyl.

1 11. The method of claim 10, wherein R³ is C₁₀ alkyl.

1 12. The method of claim 1, wherein R¹ is methyl and R² and R³ are
2 independently C₈-C₁₂ alkyl.

- 1 13. The method of claim 12, wherein R² and R³ are C₁₀ alkyl.
- 1 14. The method of claim 1, wherein R¹ and R² are methyl and R³ is C₈-C₂₀
2 alkyl.
- 1 15. The method of claim 1, wherein the amine is selected from the group
2 consisting of didecylmethanamine, dodecylmethanamine, dioctylmethanamine,
3 octadecylmethanamine, dioctadecylmethanamine, trioctylamine, and any combination of
4 any of the foregoing.
- 1 16. The method of claim 1, wherein R⁴ is hydrogen or methyl.
- 1 17. The method of claim 16, wherein the cyclic carbonate is ethylene carbonate.
- 1 18. The method of claim 16, wherein the cyclic carbonate is propylene
2 carbonate.
- 1 19. The method of claim 1, wherein R⁵, R⁶, R⁷, and R⁸ are independently
2 hydrogen or C₁-C₄ alkyl.
- 1 20. The method of claim 19, wherein R⁵, R⁶, R⁷, and R⁸ are independently
2 hydrogen or methyl.
- 1 21. The method of claim 1, wherein R⁵ is methyl and R⁶, R⁷, and R⁸ are
2 hydrogen.
- 1 22. The method of claim 1, wherein m ranges from 1 to 100.
- 1 23. The method of claim 1, wherein the molar ratio of amine to component (b)
2 ranges from about 1:1 to about 1:10.

1 24. The method of claim 23, wherein the molar ratio of amine to component (b)
2 ranges from about 1:2 to about 1:3.

1 25. The method of claim 1, wherein the molar ratio of amine to methanol
2 ranges from about 1:2 to about 1:20.

1 26. The method of claim 1, wherein the reaction step is performed at from
2 about 120 to about 160° C.

1 27. The method of claim 26, wherein the reaction step is performed at from
2 about 120 to about 150° C.

1 28. The method of claim 27, wherein the reaction step is performed at from
2 about 120 to about 140° C.

1 29. The method of claim 1, further comprising the step of recovering the
2 dimethyl carbonate.

1 30. The method of claim 1, wherein the reaction step comprises reacting
2 (a) the amine;
3 (b) (i) the cyclic carbonate,
4 (ii) the aliphatic polyester, or
5 (iii) a mixture thereof;
6 (c) methanol; and
7 (d) dimethylcarbonate.

1 31. The method of claim 30, wherein the molar ratio of amine to
2 dimethylcarbonate ranges from about 2:1 to about 1:3.


1 32. A method of preparing didecyldimethyl ammonium methocarbonate
2 comprising reacting

- 1 (a) didecylmethylamine;
 2 (b) a cyclic carbonate selected from the group consisting of ethylene carbonate,
 3 propylene carbonate, and mixtures thereof; and
 4 (c) methanol
 5 to form didecyltrimethyl ammonium methocarbonate.

- 1 33. The method of claim 32, wherein the reaction step comprises reacting
 2 (a) didecylmethylamine;
 3 (b) a cyclic carbonate selected from the group consisting of ethylene carbonate,
 4 propylene carbonate, and mixtures thereof;
 5 (c) methanol; and
 6 (d) dimethylcarbonate.

- 1 34. The method of claim 32, wherein the cyclic carbonate is propylene
 2 carbonate.

- 1 35. A method of preparing a quaternary ammonium bicarbonate having the
 2 formula



- 3
 4
 5
 6
 7
 8
 9 wherein R^1 , R^2 , and R^3 are independently C_1 - C_{30} alkyl, the method comprising
 10 (a) preparing a quaternary ammonium methocarbonate by the method of claim
 11 1; and
 12 (b) converting the quaternary ammonium methocarbonate to the quaternary
 13 ammonium bicarbonate.

1 36. A method of preparing didecyldimethyl ammonium bicarbonate comprising
2 (a) reacting
3 (i) didecylmethylamine,
4 (ii) a cyclic carbonate selected from the group consisting of ethylene
5 carbonate, propylene carbonate, and mixtures thereof, and
6 (iii) methanol
7 to form didecyldimethyl ammonium methocarbonate; and
8 (b) converting the didecyldimethyl ammonium methocarbonate to
9 didecyldimethyl ammonium bicarbonate.

1 37. The method of claim 36, wherein step (a) comprises reacting
2 (i) didecylmethylamine,
3 (ii) a cyclic carbonate selected from the group consisting of ethylene
4 carbonate, propylene carbonate, and mixtures thereof,
5 (iii) methanol, and
6 (iv) dimethylcarbonate
7 to form didecyldimethyl ammonium methocarbonate.

1 38. A method of preparing a mixture of quaternary ammonium bicarbonate and
2 quaternary ammonium carbonate wherein the quaternary ammonium cation has the
3 formula $N^+(CH_3)R^1R^2R^3$ and R^1 , R^2 , and R^3 are independently C_1 - C_{30} alkyl, the method
4 comprising
5 (a) preparing a quaternary ammonium methocarbonate by the method of claim
6 1; and
7 (b) converting the quaternary ammonium methocarbonate to a mixture of
8 quaternary ammonium bicarbonate and quaternary ammonium carbonate.

1 44. The method of claim 39, wherein the molar ratio of amine to methanol
2 ranges from about 1:2 to about 1:20.

1 45. The method of claim 39, wherein the reaction step is performed at from
2 about 120 to about 160° C.


1 46. The method of claim 45, wherein the reaction step is performed at from
2 about 120 to about 150° C.

1 47. The method of claim 46, wherein the reaction step is performed at from
2 about 120 to about 140° C.

1 48. The method of claim 39, further comprising the step of recovering alkanol
2 having the formula R^9OH .

1 49. The method of claim 39, wherein the reaction step comprises reacting
2 (a) the amine;
3 (b) the ester;
4 (c) methanol; and
5 (d) alkyl methyl carbonate having the formula $CH_3OC(O)OR^9$.


1 50. A method of preparing a quaternary ammonium bicarbonate having the
2 formula



8 wherein R^1 , R^2 , and R^3 are independently C_1 - C_{30} alkyl, the method comprising
9 (a) preparing a quaternary ammonium alkylcarbonate by the method of claim
10 39; and


- 1 (b) converting the quaternary ammonium alkylcarbonate to the quaternary
2 ammonium bicarbonate.

- 1 51. A method of preparing a quaternary ammonium methocarbonate having the
2 formula



- 3
4
5
6
7
8
9 wherein R^1 and R^2 are independently C_1 - C_{30} alkyl and R^3 is a C_8 - C_{30} alkyl, the method
10 comprising reacting

- 11 (a) an amine having the formula $NR^1R^2R^3$;
12 (b) (i) a cyclic carbonate having the formula



- 13
14
15
16
17
18
19
20 wherein R^4 is hydrogen or C_1 - C_4 alkyl and n is an integer from 1 to 10,


- 21 (ii) a polycarbonate,
22 (iii) a carbonate ester, or
23 (iv) a mixture thereof; and
24 (c) methanol
25 to form the methocarbonate.

1 52. The method of claim 51, wherein the carbonate ester has the formula

2

3

4



5 wherein R^9 is $-\text{CH}_3$ and R^{10} is a C_1 - C_{10} alkyl.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 02/21236

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C07C209/20 C07C211/63

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, BEILSTEIN Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	YAGI, O.; SHIMIZU, S.: "Synthesis of pure tetramethylammonium hydroxide solution free from chloride ion by the electrolysis of its hydrogen carbonate" CHEMISTRY LETTERS, vol. 12, 1993, pages 2041-2044, XP002218995 page 2042; table 1	1-38,51, 52
X	---	39-50
Y	EP 0 291 074 A (MITSUBISHI PETROCHEMICAL CO) 17 November 1988 (1988-11-17) page 4, line 14-24; claims 1,5; example 1	1-38,51, 52
X	---	39-50
	--- -/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the International filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the International filing date but later than the priority date claimed

T later document published after the International filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the International search

31 October 2002

Date of mailing of the International search report

27/11/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel (+31-70) 340-2040, Tx. 31 651 epo nl
Fax (+31-70) 340-3016

Authorized officer

Herzog, A

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 02/21236

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 4 062 884 A (MELIS UGO ET AL) 13 December 1977 (1977-12-13) cited in the application the whole document ---	1-38, 51, 52
X	US 2 635 100 A (WERNITZ JAMES H) 14 April 1953 (1953-04-14) cited in the application example V ---	39, 41, 42, 44, 49
A	DATABASE WPI Section Ch, Week 199141 Derwent Publications Ltd., London, GB; Class E14, AN 1991-298749 XP002218996 & JP 03 197449 A (MITSUI PETROCHEM IND CO LTD), 28 August 1991 (1991-08-28) abstract ---	1-52
A	DATABASE WPI Section Ch, Week 199517 Derwent Publications Ltd., London, GB; Class B05, AN 1995-128295 XP002218997 & JP 07 053478 A (MITSUBISHI GAS CHEM CO INC), 28 February 1995 (1995-02-28) abstract ---	1-52
A	US 5 523 487 A (WALKER LEIGH E) 4 June 1996 (1996-06-04) cited in the application abstract; claims; examples -----	1-52

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 02/21236

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 0291074	A	17-11-1988	JP 8019060 B	28-02-1996
			JP 63280045 A	17-11-1988
			JP 7116113 B	13-12-1995
			JP 63284148 A	21-11-1988
			JP 2114597 C	06-12-1996
			JP 8026052 B	13-03-1996
			JP 63290889 A	28-11-1988
			JP 2114598 C	06-12-1996
			JP 8026053 B	13-03-1996
			JP 63290890 A	28-11-1988
			DE 3868138 D1	12-03-1992
			EP 0291074 A2	17-11-1988
			US 4892944 A	09-01-1990
US 4062884	A	13-12-1977	IT 1034961 B	10-10-1979
			CA 1064047 A1	09-10-1979
			DE 2615665 A1	14-10-1976
			DK 145776 A ,B,	10-10-1976
			FR 2306972 A1	05-11-1976
			GB 1489736 A	26-10-1977
			IE 43223 B1	14-01-1981
			JP 51122025 A	25-10-1976
			JP 59028542 B	13-07-1984
			LU 74710 A1	11-11-1976
			NL 7603679 A ,B	12-10-1976
			NO 761213 A ,B,	12-10-1976
US 2635100	A	14-04-1953	NONE	
JP 3197449	A	28-08-1991	NONE	
JP 7053478	A	28-02-1995	NONE	
US 5523487	A	04-06-1996	US 5438034 A	01-08-1995
			US 6090855 A	18-07-2000
			AU 7107694 A	03-01-1995
			CA 2162128 A1	22-12-1994
			CA 2315849 A1	22-12-1994
			EP 1114704 A2	11-07-2001
			EP 1122044 A1	08-08-2001
			EP 1121857 A1	08-08-2001
			EP 0702517 A1	27-03-1996
			IL 109964 A	22-09-1999
			IL 122727 A	12-03-1999
			IL 122728 A	12-03-1999
			IL 122729 A	12-03-1999
			IL 122730 A	12-03-1999
			IL 124041 A	31-01-2000
			JP 8511543 T	03-12-1996
			NZ 268249 A	26-08-1998
			WO 9428715 A1	22-12-1994
			ZA 9403999 A	03-02-1995